



**City of Philadelphia
Department of Public Health
Air Management Services**

Philadelphia's Air Quality Report 2008





City of Philadelphia
Department of Public Health
Air Management Services

Executive Summary

This report focuses on the air quality of the City of Philadelphia, as presented by the Philadelphia Department of Public Health, Air Management Services (AMS), the local air pollution control agency for the City of Philadelphia. As an urban area, Philadelphia faces many of the same pollution challenges as other densely populated areas, such as emissions from vehicles and industries. The information contained in this report reviews Philadelphia's air quality for the year 2008, and reports how the City's air compared with the National Ambient Air Quality Standards (NAAQS). This report covers the following criteria pollutants: **ozone, carbon monoxide, nitrogen dioxide, sulfur dioxide, particulate matter, and lead**. It also provides an overview of **Hazardous Air Pollutants** also referred to as air toxics. There are no national standards for air toxics.

In general, trends show many air pollutants in Philadelphia to be decreasing. In 2008, Philadelphia was considered to be in attainment for all pollutants, with the exception of ozone and particulate matter that is less than 2.5 microns in diameter. There were 204 good days, 144 moderate days, and 18 unhealthy days in Philadelphia. In addition, air toxics that were measured in Philadelphia that show an excess lifetime cancer risk of more than one in a million were: 1,3-butadiene, acetaldehyde, benzene, carbon tetrachloride and tetrachloroethylene.

For further information, please visit the Air Management Services website at:

<http://www.phila.gov/health/units/ams/index.html>

or contact us at:

215-685-7580

During non-business hours, call 311. Operators take calls 8 a.m. to 8 p.m. on weekdays and 9 a.m. to 5 p.m. on Saturdays. It is closed on Sundays.

Michael A. Nutter, Mayor
Donald F. Schwarz, Deputy Mayor for Health and Opportunity and Health
Commissioner



Table of Contents

Introduction	5
Air Monitoring Network	5
Quality Assurance	7
Air Quality Index	8
National Ambient Air Quality Standards	11
The Pollutants We Measure	11
Ozone (O ₃)	12
Carbon Monoxide (CO)	15
Nitrogen Dioxide (NO ₂)	17
Sulfur Dioxide (SO ₂)	18
Lead (Pb)	20
Particulate Matter (PM ₁₀ , PM _{2.5})	22
PM ₁₀	23
PM _{2.5}	25
Air Toxics	27
Appendices	
Appendix A: Consequences of Air Pollution	37
Appendix B: What AMS is Doing to Reduce Air Pollution	38
Appendix C: Protecting Yourself and the Environment	39
Appendix D: Frequently Asked Questions About Air Quality	40
Appendix E: Websites	43
Appendix F: Contacts	44
Figures	
Figure 1 - 2008 Philadelphia Air Monitoring Network	6
Figure 2 - Color Coded Air Quality Index (AQI)	9
Figure 3 - Philadelphia Annual AQI Summary Standardized by Current EPA Breakpoints	10
Figure 4.1 - Ozone Monitoring Map	12
Figure 4.2 - Ozone Trends (Minimum, Maximum, Average) for Highest 4 th Daily Maximum 8-Hour Concentration	14
Figure 4.3 - Ozone 8-Hour Design Value for NEA Monitoring Site (421010024)	14
Figure 5.1 - CO Monitoring Map	15
Figure 5.2 - CO Trends (Minimum, Maximum, Average) for Highest 2 nd Maximum 8- Hour Concentration	16



Figure 6.1 - NO ₂ Monitoring Map.....	17
Figure 6.2 - NO ₂ Trends (Minimum, Maximum, Average) for Annual Concentration	17
Figure 7.1 - SO ₂ Monitoring Map.....	18
Figure 7.2 - SO ₂ Trends (Minimum, Maximum, Average) for Highest Annual Concentration.....	19
Figure 7.3 - SO ₂ Trends (Minimum, Maximum, Average) for Highest 2 nd Maximum 24-Hour Concentration.....	19
Figure 8.1 - Lead Monitoring Map	20
Figure 8.2 - Lead Trends (Minimum, Maximum, Average) for Highest Quarterly Maximum Concentration for Areas Near Castor and Delaware Avenues.....	21
Figure 9.1 - PM ₁₀ Monitoring Map	23
Figure 9.2 - PM ₁₀ Trends (Minimum, Maximum, Average) for Highest 2 nd Maximum 24- Hour Concentration for Areas Near Castor and Delaware Avenues	24
Figure 9.3 - PM ₁₀ Trends (Minimum, Maximum, Average) for Highest 2 nd Maximum 24- Hour Concentration for Areas Excluding Castor and Delaware Avenues.....	24
Figure 10.1 - PM _{2.5} Monitoring Map.....	25
Figure 10.2 - PM _{2.5} Annual Design Value for CHS Monitoring Site (421010047)	26
Figure 10.3 - PM _{2.5} Trends 98 th Percentile Value (Minimum, Maximum, Average) for 24- Hour Concentration	26
Figure 11 - Air Toxics Monitoring Map.....	27
Figure 12.1 - 1,3-Butadiene Average Concentration for Five Monitoring Sites.....	30
Figure 12.2 - 2002 NATA Total Cancer Risk for 1,3-Butadiene.....	30
Figure 12.3 - 1,3-Butadiene Sources of Emissions	30
Figure 13.1 - Acetaldehyde Average Concentration for Five Monitoring Sites	31
Figure 13.2 - 2002 NATA Total Cancer Risk for Acetaldehyde	31
Figure 13.3 - Acetaldehyde Sources of Emissions.....	31
Figure 14.1 - Benzene Average Concentration for Five Monitoring Site	32
Figure 14.2 - 2002 NATA Total Cancer Risk for Benzene.....	32
Figure 14.3 - Benzene Sources of Emissions	32
Figure 15.1 - Carbon Tetrachloride Average Concentration for Five Monitoring Sites ..	33
Figure 15.2 - 2002 NATA Total Cancer Risk for Carbon Tetrachloride	33
Figure 15.3 - Carbon Tetrachloride Sources of Emissions	33
Figure 16.1 - Tetrachloroethylene Average Concentration for Five Monitoring Sites	34
Figure 16.2 - 2002 NATA Total Cancer Risk for Tetrachloroethylene	34
Figure 16.3 - Tetrachloroethylene Sources of Emissions.....	34
Figure 17.1 - 2002 EPA NATA Total Cancer Risk for Philadelphia	35
Figure 17.2 - Sum of HAPs Sources of Emissions.....	35
Figure 18 - Cancer Risks for the Top Five Toxics Compounds	36



Introduction

AMS is responsible for the prevention, abatement, and control of air pollution and air pollution nuisances, achieving and maintaining federal National Ambient Air Quality Standards (NAAQS) in Philadelphia, and protecting the health and quality of life of the Philadelphia community from the adverse effects of air contaminants and noise.

AMS implements the environmental protection mandates contained in city, state, and federal laws, reviews construction and operating permits for compliance with air standards and guidelines, operates and maintains a citywide air sampling network to continuously monitor Philadelphia's air, routinely inspects pollution sources, services citywide complaints of air pollution, asbestos, and noise, issues violations, conducts enforcement actions, and advances voluntary emissions reductions.

Air Monitoring Network

The City of Philadelphia is served by a network of ten air monitoring sites located throughout the City that measure the criteria pollutants: ozone (O_3), carbon monoxide (CO), nitrogen dioxide (NO_2), sulfur dioxide (SO_2), particulate matter (PM_{10} and $PM_{2.5}$), and lead (Pb). Five of the sites also measure toxics, such as 1,3-butadiene, benzene and carbon tetrachloride. Many of the measurements are made in "real time", meaning that the measurements show pollution levels as they occur, instead of after the fact. The map on page 6 shows the location of air monitors and the pollutants measured at each monitor location. AMS measures air quality for several reasons:

- To ensure that long-term goals and targets to reduce levels of air pollution are being met.
- To provide information to the public as to how good or bad the air quality is in Philadelphia.
- To ensure attainment with standards set forth by the US EPA.

An annual air monitoring network plan has been made available to the public starting in the Year 2007 and is located on the AMS website:

<http://www.phila.gov/health/units/ams/index.html>.



Quality Assurance

The AMS Air Monitoring Laboratory's main responsibility is to provide accurate data on the quality of the City's air. Consequently, much effort is spent to achieve this objective. The measurement of pollutants in the atmosphere is being done to answer a number of questions such as:

- Are the National Ambient Air Quality Standards (NAAQS) being met in Philadelphia?
- How close or far away are we from meeting these standards?
- Which pollutants are getting worse (increasing in concentration) or better?

Many of our measurements require detecting very small amounts of a pollutant, often expressed as parts per million (ppm) or parts per billion (ppb). An illustration: imagine a million yellow balls all the same size with several red balls in the middle of them; we would need to find those red balls and then be able to count them. The instruments used to measure air pollutants need to be reliable in identifying the pollutant and accurate in making the measurement every time. The main way we check to see if our instruments are giving accurate measurements is to send a sample of air which has a known amount (concentration) of a pollutant and compare what the instrument says is the concentration to what we know is the right concentration. Then adjustments (calibration) to an instrument can be made to give a better measurement. If the needed correction is too great, the instrument needs to be repaired or replaced. EPA and our Laboratory have guidelines on how good and reliable measurements need to be to answer the questions being asked. The instruments being used now are much more reliable than those available years ago. Steps to assure good data quality include:

- Automated calibration.
- Manual calibration conducted by chemists.
- Review of the data by an experienced engineer or scientist.

The system is geared towards public safety; for example, a few measurements can be enough to identify a problem in meeting one of the NAAQS, but many good measurements over a period of time (often three years) as well as additional types of analysis are needed to "demonstrate compliance" with one of the NAAQS.



Air Quality Index

The Air Quality Index (AQI) is a color coding system for air quality used by government agencies across the United States. Media outlets disseminate air quality reports using the AQI to help warn the public about day-to-day pollution problems. Air quality alerts are issued when pollution is rated as Orange (Unhealthy for Sensitive Groups), Red (Unhealthy) or Maroon (Very Unhealthy). Alerts are more likely to occur in the summer months, but can happen any time of the year.

The AQI is needed to report pollutant levels based on five criteria air pollutants: ground-level ozone, particulate matter, carbon monoxide, sulfur dioxide, and nitrogen dioxide. Using formulas created by the EPA, data for each pollutant is converted into a score ranging from 1 to 500. Values exceeding 100, which generally correspond to the National Ambient Air Quality Standard for each pollutant, trigger "Action Days," where the public is advised to do their part to reduce pollution and take precautions to protect themselves and their families from health effects. For example, on an Orange day, children, seniors and those with respiratory ailments are advised to minimize prolonged outdoor exposure. On a Red day, all residents are advised to limit outdoor activity. The highest of the five pollutant scores is reported as the overall air quality rating for Philadelphia for a given day. That is, any individual pollutant can, on its own, trigger an Action Day. Action Days are reported through print, radio and television media and by local and regional air agencies.

Philadelphia's Real-Time Air Quality Website is located at www.phila.gov/aqi and provides the most up-to-date information about the air quality in Philadelphia. It lets you know what you should do to protect your health if the air quality is unhealthy.

Figure 2 on the following page shows the AQI summaries, as used by media outlets. The recommended actions that individuals should take to protect their health and plan their daily activities are described below the index.

Figure 2 - Color Coded Air Quality Index (AQI)

Air Quality Index (AQI) Values	Levels of Health Concern	Colors
When the AQI is in this range:	...air quality conditions are:	...as symbolized by this color:
0 to 50	Good	Green
51 to 100	Moderate	Yellow
101 to 150	Unhealthy for Sensitive Groups	Orange
151 to 200	Unhealthy	Red
201 to 300	Very Unhealthy	Purple
301 to 500	Hazardous	Maroon

"Good" The AQI value is between 0 and 50. Air quality is considered satisfactory, and air pollution poses little or no risk.

"Moderate" The AQI is between 51 and 100. Air quality is acceptable; however, for some pollutants there may be a moderate health concern for a very small number of people. For example, people who are unusually sensitive to ozone may experience respiratory symptoms.

"Unhealthy for Sensitive Groups" When AQI values are between 101 and 150, members of sensitive groups may experience health effects. This means they are likely to be affected at lower levels than the general public. For example, people with lung disease are at greater risk from exposure to ozone, while people with either lung disease or heart disease are at greater risk from exposure to particle pollution. The general public is not likely to be affected when the AQI is in this range.

"Unhealthy" Everyone may begin to experience health effects when AQI values are between 151 and 200. Members of sensitive groups may experience more serious health effects.

"Very Unhealthy" AQI values between 201 and 300 trigger a health alert, meaning everyone may experience more serious health effects.

"Hazardous" AQI values over 300 trigger health warnings of emergency conditions. The entire population is more likely to be affected.



City of Philadelphia
 Department of Public Health
 Air Management Services

Figure 3 below shows the annual summary of the number of good, moderate, and unhealthy air quality days in Philadelphia based on monitoring conducted by AMS since 1990.

Figure 3 - Philadelphia Annual AQI Summary Standardized by Current EPA Breakpoints



The number of days with good air quality steadily increased until 1999 and then decreased until 2002 before again increasing and subsequently leveling off. In the same timeframe, the number of days with moderate air quality increased and leveled off. Over the period from 1990 through 1998, the annual number of days with unhealthy air quality dramatically decreased and has remained about the same each year since. These improvements can be attributed mainly to emission reductions from gasoline marketing, including vapor recovery at retail gasoline stations, and companies shutting down pollution producing processes.

Figure 3 has been standardized with the current AQI breakpoints. Stricter standards have come online including 1997 national standards for ozone and PM_{2.5}. The 8-hour standard for ozone was revised in 2008 to 0.075 ppm from 0.08 ppm and the 24-hour standard for PM_{2.5} was revised in 2006 to 35 µg/m³ from 65 µg/m³. As a result of more stringent standards, some days that in prior years would have been reported as good fell into the moderate range and those that would have been reported as moderate fell



into the unhealthy range. Therefore, Figure 3 does not reflect real deterioration in air quality since 1999. The air quality in Philadelphia is actually getting better.

National Ambient Air Quality Standards

The Clean Air Act of 1970 empowered EPA to establish National Ambient Air Quality Standards (NAAQS) to protect public health and welfare. Since that time, standards have been set for: carbon monoxide, sulfur dioxide, nitrogen dioxide, lead, particulate matter (“dust” or “soot”) and ozone. The NAAQS is expressed as a pollutant concentration averaged over a fixed amount of time. For example, the NAAQS for sulfur dioxide are:

- 0.5 parts of sulfur dioxide per million parts of air over a time period of 3-hours.
- 0.14 parts of sulfur dioxide per million parts of air over a time period of 24-hours.
- 0.03 parts of sulfur dioxide per million parts of air over a time period of one year.

In 2008, Philadelphia was considered to be in attainment for all pollutants, with the exception of ozone and particulate matter that is less than 2.5 microns in diameter.

The Pollutants We Measure

The following sections provide information on the health effects, sources and trends of pollutants measured in Philadelphia. Included are the pollutants that EPA established National Ambient Air Quality Standards (NAAQS), commonly called criteria pollutants, as well as pollutants identified as being toxic or hazardous. Each of the criteria pollutants are graphed to show the historical trend compared with national standards. The graphs also identify the sites of the “worst” levels, the “best” levels, and with a solid circle, the mean level (the average of all recorded levels). It is important to note the mean, as it factors out extreme levels, and thereby provides a better indication of the average for air quality.




Ozone (O₃)

NAAQS:

1997: Highest 4th Daily Maximum 8-Hour Concentration = 0.08 ppm

2008: Highest 4th Daily Maximum 8-Hour Concentration = 0.075 ppm

Figure 4.1 - Ozone Monitoring Map



Ground level ozone (the primary constituent of smog) is the pollutant most often responsible for unhealthy air quality in the Philadelphia region. Ozone is not emitted into the atmosphere directly but is formed by chemical reactions between other pollutants. Volatile Organic Compounds (VOCs) and Nitrogen Oxides (NO_x) react to create ozone in the presence of heat and sunlight. Ozone levels are consistently higher during the summer months.

There are four categories of emission sources from human activity that produce VOC and NO_x. The four categories are: **Point sources** - the largest utilities, industries, and other operations; **Area Sources** - commercial, solvent use, waste disposal, and other categories; **Non-road Engine Sources** - construction and agricultural equipment,

recreational boats, and lawnmowers; **Highway Vehicle Sources** - cars, trucks, buses, and motorcycles. Emissions of VOC and NO_x may be carried by wind currents while reacting to produce high ozone levels hundreds of miles from their sources. In the eastern United States during the summer months, ground level ozone is frequently high over wide areas containing several states, caused by ozone transfer.

Unlike the oxygen that we breathe, which has only two atoms of oxygen (O₂), ozone has an additional oxygen atom, making it very reactive. This is why ozone is said to burn or irritate the lungs. People who are very young or very old, or who have chronic lung problems such as asthma are particularly sensitive to ground level ozone.

In any discussion of ozone, it is important to distinguish between the effects of ozone at the ground and ozone high in the atmosphere, several miles above our heads. An advertisement might use the slogan "good up high, bad nearby," to describe ozone. Regardless of where it is, no one would want to breathe it. However, up high in what's called the ozone layer, ozone is essential to the health of nearly every living thing, since it protects the Earth from harmful ultraviolet (UV) light. If not for this natural layer, UV light would sterilize the Earth's surface, and life as we know it would cease to exist.



City of Philadelphia
Department of Public Health
Air Management Services

Near the ground, ozone reacts with buildings, plants, animals, and people, and is one of the most irritating, harmful components of smog. Smog refers to the whole mixture of air pollution in an area, and may include ozone, a whole host of other gases, as well as fine particles and the hazy conditions they cause.

VOCs are organic compounds that evaporate readily, such as gasoline vapors and paint fumes. NO_x stands for two compounds, nitric oxide (NO) and nitrogen dioxide (NO₂). VOCs that come from human activities are called anthropogenic VOCs. Some anthropogenic VOCs, such as benzene, are themselves toxic and may increase risks of cancer or lead to other adverse health effects in addition to helping form ozone. Some VOCs are considerably more reactive in the atmosphere than others, and the reactivity of a VOC influences how quickly ozone forms. A compound that reacts in a few minutes to produce ozone will have a much greater impact near its source than one that reacts more slowly. Thus, ozone can form at various distances downwind of a VOC source due to the speed of these chemical reactions.

On July 18, 1997, the EPA promulgated a revision to the National Ambient Air Quality Standard for ozone which came into effect on July 1, 1999. Previously, the standard was based on the number of times that the daily maximum hourly ozone concentration was greater than 0.12 parts per million (ppm) over a 3-year period. The revised NAAQS is based on an 8-hour average ozone concentration. EPA revoked the 1-hour standard on June 15, 2005. The latest ozone standard of 0.075 ppm for the 8-hour average ozone concentration has been effective since May 27, 2008. This standard replaced the 1997 standard of 0.08 ppm.

In 2008, Philadelphia and the surrounding counties were in nonattainment for the 8-hour ozone standard. This means that the standard set by the EPA for ozone was exceeded. This standard was exceeded 15 times in 2008. AMS, along with other local and regional air quality agencies, continues to work towards compliance with ozone standards.

A State Implementation Plan (SIP) is a plan which identifies how a state will attain the standard. Each state is required to have a SIP which contains control measures and strategies that demonstrate how each area will attain and maintain the NAAQS. These plans are developed through a public process, formally adopted by the state, and submitted by the Governor's designee to the EPA.

Starting in 2008, ELM and ROX monitoring sites were no longer active in the ozone monitoring network. Currently, as seen in Figure 4.1, there are two active ozone monitoring sites: LAB and NEA.

Figure 4.2, on the following page, shows the trends for the ozone 8-hour concentration in Philadelphia.



Figure 4.2 - Ozone Trends (Minimum, Maximum, Average) for Highest 4th Daily Maximum 8-Hour Concentration





Figure 4.3 - Ozone 8-Hour Design Value for NEA Monitoring Site (421010024)






Carbon Monoxide (CO)

NAAQS:

Highest 2nd Maximum 8-Hour Concentration = 9 ppm

Figure 5.1 - CO Monitoring Map



Carbon monoxide (CO) is colorless, odorless, and at high concentrations is a poisonous gas. It is formed when carbon in fuels are not burned completely. The major source of CO is motor vehicle emissions. Other sources of CO include residential, industrial, and natural processes. Weather greatly affects CO levels, and peak CO concentrations typically occur during the colder months of the year.

Carbon monoxide enters the bloodstream and reduces oxygen delivery to the body's organs and tissues. The health threat from carbon monoxide is most serious for those who suffer from cardiovascular disease. Exposure to elevated CO levels is associated with impairment of vision, reduced work capacity, reduced manual dexterity,


poor learning ability, and difficulty in performing complex tasks. At very high levels, carbon monoxide can be fatal.

Over a nineteen year period, there has been a continued reduction in carbon monoxide levels. This is mainly the result of federal requirements for cleaner automobiles and fuel and state inspection/maintenance programs. NBR and CHS monitoring sites are no longer active in the CO monitoring network. The LAB is operating as the main monitoring site for CO.

Figure 5.2, on the following page, shows the trends for the CO 8-hour concentration in Philadelphia.



Figure 5.2 - CO Trends (Minimum, Maximum, Average) for Highest 2nd Maximum 8-Hour Concentration






Nitrogen Dioxide (NO₂)

NAAQS:

Highest Annual Arithmetic Mean Concentration = 0.053 ppm

Figure 6.1 - NO₂ Monitoring Map




Nitrogen dioxide is a light brown gas that is an important component of urban haze. The compound is created primarily from fuel combustion in motor vehicles, utilities, and industrial sources.

Nitrogen dioxide can irritate the lungs and lower resistance to respiratory infections such as influenza. Nitrogen oxides (NO_x) are an important precursor to both ozone and acid rain and can affect both land and water ecosystems. They contribute to the formation of fine particulate matter, haze and reductions in visibility.

Ambient levels of nitrogen dioxide in Philadelphia are better than the NAAQS showing a sustained downward trend over time.

Figure 6.2 - NO₂ Trends (Minimum, Maximum, Average) for Annual Concentration






Sulfur Dioxide (SO₂)

NAAQS:

Highest Annual Mean Concentration = 0.03 ppm

Highest 2nd Maximum 24-Hour Concentration = 0.14 ppm

Figure 7.1 - SO₂ Monitoring Map



SO₂ is emitted from the burning of fuels that contain sulfur. Industrial grade fuel oils are the primary source in Philadelphia.

The major health concerns associated with exposure to high concentrations of SO₂ include effects on breathing, respiratory illness, alterations in the lungs' defenses, and aggravation of existing respiratory and cardiovascular disease. Together, SO₂ and NO_x are the major ingredients of acid rain. SO₂ also plays a significant role in the formation of fine particulate matter.

SO₂ levels are well within air quality standards and show a slow, continued improvement over time. This is mainly due to industry, businesses, and

homes changing to fuels with lower sulfur content such as natural gas.

AMS has shutdown the operation of the ELM site as one of the SO₂ monitoring sites in the network; LAB and RIT sites are still operating as seen in Figure 7.1

The following graphs, Figure 7.2 and Figure 7.3 show the trends for the SO₂ annual mean concentration and 24-hour concentration, respectively for Philadelphia.



Figure 7.2 - SO₂ Trends (Minimum, Maximum, Average) for Highest Annual Concentration





Figure 7.3 - SO₂ Trends (Minimum, Maximum, Average) for Highest 2nd Maximum 24-Hour Concentration






Lead (Pb)

NAAQS:

1978: Quarterly Average Concentration = 1.5 $\mu\text{g}/\text{m}^3$

2008: Rolling 3-Month Average Concentration = 0.15 $\mu\text{g}/\text{m}^3$

Figure 8.1 - Lead Monitoring Map



The processing of metals is the major source of lead emissions to the atmosphere. Lead does not travel over great distances in the air and so concentrations vary, with highest levels near specific industrial sites.

Lead is a metal that is highly toxic when inhaled or ingested. Lead accumulates in the blood, bone, and soft tissue and may affect the kidneys, liver, nervous system and other organs. It also can cause learning difficulties in children.

Ambient lead levels have been decreasing throughout the city due to the elimination of leaded gasoline and greater control of emissions from companies that produce or process lead

compounds.

Lead levels in certain parts of the city were once extremely high due to the concentration of particular industries in the area, and this is reflected in the high readings for monitors near Castor and Delaware Avenues before 1998. The levels of lead in these areas have drastically improved, and are now comparable to the rest of the city.


On October 15, 2008, the EPA strengthened its regulation for lead. The standard was revised from the 1978 standard of 1.5 $\mu\text{g}/\text{m}^3$ to a level that is 10 times more stringent, 0.15 $\mu\text{g}/\text{m}^3$, with a different averaging time. For the previous standard, the averaging time used a quarterly average while the new standard uses a rolling 3-month average. The revision is based on more than 6000 studies performed since 1990 on the health effects of high lead concentrations in the bloodstream. The studies show that adverse effects from lead in the blood occur at a much lower level than previously thought.

There is no change in the status of attainment for Philadelphia regarding the new regulation. Currently, AMS measures for ambient lead at the historically highest location, ITO.



City of Philadelphia
Department of Public Health
Air Management Services

Figure 8.2 - Lead Trends (Minimum, Maximum, Average) for Highest Quarterly Maximum Concentration for Areas Near Castor and Delaware Avenues





Particulate Matter (PM₁₀, PM_{2.5})

Particulate matter is the general term used for a mixture of solid particles and liquid droplets found in the air. These particles come in a wide range of sizes and originate from stationary, mobile, and natural sources.

PM₁₀ and PM_{2.5} refer to small particulates that measure less than 10 micrometers (0.00001 meters) and 2.5 micrometers (0.0000025 meters) in diameter, respectively. In addition to health problems, particulate matter can cause reduced visibility, soiling, and damage to materials. Particles of this size remain airborne for long periods of time and disperse in uniform concentrations across wide areas, crossing geographic boundaries.

In 1997, the EPA set a separate standard for PM_{2.5}. Particles in the PM_{2.5} size range are able to travel deeply into the respiratory tract, reaching the lungs. Exposure to fine particles can cause short-term health effects such as eye, nose, throat and lung irritation, coughing, sneezing, runny nose and shortness of breath. Exposure to fine particles can also affect lung function and worsen medical conditions such as asthma and heart disease. Scientific studies have linked increases in daily PM_{2.5} exposure with increased respiratory and cardiovascular hospital admissions, emergency department visits and deaths. Recent studies suggest that long term exposure to particulate matter may be associated with increased rates of bronchitis and reduced lung function.

Particles come in a wide variety of shapes and sizes, which affect their impacts on the environment and human health. Bigger particles, such as dust, are easier to see and can cause problems, but smaller particles are likely to be worse for our health.

Fine particles are treated as though they are a single pollutant, but fine particles come from many different sources and are composed of thousands of different compounds. Fortunately, these compounds fall into a few dominant categories: sulfates, nitrates, ammonium compounds, soil, organic carbon compounds, and elemental carbon. Water is nearly always an important and variable part of PM, and sea salt is often significant near the coast. Given the complex composition of PM, it is no surprise that its chemistry is also complex. Particles may be dry or wet. When the wind blows hard enough, soil, silt, and sand can be lifted from the surface. Human activities such as mining, construction, plowing, and driving on unpaved roads, also lift particles into the air. Soot, also referred to as black carbon or elemental carbon, is emitted directly by diesel engines and forest fires, among other sources. Most individual particles are likely mixtures of different substances, the products of growing by collisions with other particles and by taking on gases.




PM₁₀

NAAQS:

Highest 2nd Maximum 24-Hour Concentration = 150 µg/m³ (not to be exceeded more than once per year on average over 3 years)

Figure 9.1 - PM₁₀ Monitoring Map



Particulate matter levels have been decreasing due to regulations limiting the amount of emissions allowed and the change to cleaner fuels such as natural gas by industry, businesses and homes.

During the mid 1990s, particulate emissions from several sources in the area of Castor and Delaware Avenues caused extremely high localized measurements. In fact, the levels were many times higher than those measured at other city locations. Because the impact was not widespread, the additional charts are presented to highlight that fact. Specific actions to abate these sources have resulted in air quality that now meets the national standards and are now comparable to levels in the rest of the city.

The EPA has revoked the annual standard for PM₁₀ since December 17, 2006 due to a lack of evidence linking health problems to long-term exposure to coarse particle pollution.

ELM and VET monitoring sites no longer measure PM₁₀ in the monitoring network. The active monitoring sites as seen in Figure 9.1 are LAB, NEW, NEL and ITO.

Prior to 1998, a facility near Castor and Delaware Avenues called the Franklin Smelting and Refining Company generated significant dust from smelting activities. This facility has since shut down. The shutdown has increased the quality of the air in the area near the facility since that time.



Figure 9.2 - PM₁₀ Trends (Minimum, Maximum, Average) for Highest 2nd Maximum 24-Hour Concentration for Areas Near Castor and Delaware Avenues





Figure 9.3 - PM₁₀ Trends (Minimum, Maximum, Average) for Highest 2nd Maximum 24-Hour Concentration for Areas Excluding Castor and Delaware Avenues






PM_{2.5}

NAAQS:

Highest Annual Mean Concentration = 15 $\mu\text{g}/\text{m}^3$

Highest 98th Percentile 24-Hour Concentration = 35 $\mu\text{g}/\text{m}^3$

Figure 10.1 - PM_{2.5} Monitoring Map



PM_{2.5} consists of those particles that are less than 2.5 micrometers in diameter. They are also referred to as "fine" particles. Fine particles result from fuel combustion from motor vehicles, power generation, and industrial facilities, as well as from residential fireplaces and wood stoves. A significant amount of fine particles are also formed in the atmosphere by the transformation of gaseous emissions such as SO₂, NO_x, VOCs, and ammonia.

Fine particles can accumulate in the respiratory system and are associated with numerous health effects such as premature death, increased respiratory symptoms and disease, and decreased lung function. Sensitive groups that appear to be at

the greatest risk for such effects include the elderly, children, and individuals with cardiopulmonary disease or respiratory ailments such as asthma.

Revisions to the primary (health-based) NAAQS added two new PM_{2.5} standards, set at 15 $\mu\text{g}/\text{m}^3$ (annual standard) and at 65 $\mu\text{g}/\text{m}^3$ (24-hour standard). On December 18, 2006 the 24-hour standard was strengthened to 35 $\mu\text{g}/\text{m}^3$. Measuring PM_{2.5} requires highly sensitive equipment under tight temperature and humidity control. The charts on the next page show that the Philadelphia region is in nonattainment for the 24-hour standard of 35 $\mu\text{g}/\text{m}^3$ and is in attainment for the annual PM_{2.5} standard. The design value is defined as the highest weighted annual average concentration at any monitoring site in Philadelphia. The 24-hour concentration is the highest concentration of 98th percentile values at any monitoring site in Philadelphia.

There were some changes to the PM_{2.5} monitoring network in 2008. Two new monitors at FAB and RIT became operational while ELM and EWK sites shut down. The new FAB monitoring site located at 3rd and Spring Garden Streets, was established as an alternative to the CHS monitoring site which may be shut down in the near future. The FAB site was selected to monitor PM_{2.5} because it is in a high population area and is close to Interstate 95. An additional PM_{2.5} monitor at the RIT site was established to study the impact of the Sunoco petroleum refinery on the local community. Currently, there are five PM_{2.5} monitoring sites in the network (Figure 10.1).



Designations for the PM_{2.5} nonattainment area will not be determined until September 2009. Figure 10.2 is the design value for the CHS monitoring site. Be advised that the data may be below 75% data capture.

Figure 10.2 - PM_{2.5} Annual Design Value for CHS Monitoring Site (421010047)






Figure 10.3 - PM_{2.5} Trends 98th Percentile Value (Minimum, Maximum, Average) for 24-Hour Concentration





Air Toxics

Figure 11 - Air Toxics Monitoring Map



Air toxics, also referred to as toxic air pollutants or hazardous air pollutants (HAPs), are substances that cause adverse health effects or environmental damage. The Federal Clean Air Act Amendments (CAAA) of 1990 list 187 pollutants or chemical groups as HAPs. Examples of air toxics include heavy metals (such as beryllium), organic chemicals (such as formaldehyde), polycyclic organic matter (POM, which are formed primarily by combustion), benzene (which is found in gasoline), pesticides, fine mineral fibers, and asbestos. HAPs are emitted from stationary sources (large industrial facilities), area sources (dry cleaners and household uses), as well as mobile sources (trucks and buses).

There is less information known about the health impact from the 187 HAPs than there are for criteria pollutants, and no national standards exist for them. However, a number of these pollutants are known or suspected to be carcinogenic, and there is no known “safe concentration.” The danger posed by toxics is often referred to in terms of risk. Risk is defined as the likelihood of a negative outcome from a certain level of a specific chemical, or the measure of a chance that health problems will occur. For example, many toxics cause cancer, while others cause respiratory problems, birth defects, neurological or, immune response problems, and other health concerns. Toxics have varying degrees of danger, and some will cause harm with a very small amount of the substance while others require large amounts to have a negative effect. A risk level of one in a million implies a likelihood that up to one person, out of one million equally exposed people would contract cancer if exposed continuously (24 hours per day) to the specific concentration over 70 years (an assumed lifetime). This risk is calculated as additional to those cancer cases that would normally occur in an unexposed population of one million people.

AMS is helping to reduce HAPs in Philadelphia by enforcing Federal, State, and locally mandated programs that limit emissions from stationary and area sources. Many toxic emissions have been reduced by regulations designed to bring Philadelphia into compliance with the NAAQS for Ozone. In addition, Philadelphia enforces the National Emission Standards for Hazardous Air Pollutants (NESHAPs), a program designed to reduce emissions from existing major and area sources, as well as New Source Performance Standards (NSPS), which limit toxic emissions from new sources.



City of Philadelphia
Department of Public Health
Air Management Services

Since diesel emissions are a significant but unquantified contributing factor in determining health risks from toxic emissions, AMS continues working to promote voluntary emissions reductions from diesel vehicles and to bring clean diesel technology to the Philadelphia area. The Philadelphia Diesel Difference Working Group, a coalition of diverse stakeholders whose primary purpose is to reduce the air pollutants associated with diesel-powered engines in the greater Philadelphia area, meets on a monthly basis. More information on this program can be found at www.cleanair.org/dieseldifference.

AMS has historically measured toxic pollutants at the Laboratory (LAB) and more recently at the Community Health Services (CHS), Elmwood (ELM), Roxborough (ROX), and Ritner (RIT) monitoring sites as seen on page 27.

As part of EPA's National Air Toxics Assessment (NATA) activities, 180 air pollutants were assessed for either lifetime cancer risk or non-cancer hazard due to inhalation. NATA is EPA's ongoing comprehensive evaluation of air toxics in the U.S. These activities include: expansion of air toxics monitoring, improving and periodically updating emission inventories, improving national- and local-scale modeling, continued research on health effects and exposures to both ambient and indoor air, and improvement of assessment tools.

The goal of NATA is to identify those air toxics which are of greatest potential concern, in terms of contribution to population risk. The results are used to establish strategies to reduce emissions and these set priorities or programs and the collection of additional air toxics data.

The assessment includes four steps that look at the year 2002, the latest year with a published NATA report:

1. Compiling a national emissions inventory of air toxics emissions from outdoor sources.
2. Estimating ambient concentrations of air toxics across the contiguous United States.
3. Estimating population exposures across the contiguous United States.
4. Characterizing potential public health risk due to inhalation of air toxics including both cancer and non-cancer effects.

The 2002 NATA identified certain health risks in the City. Philadelphia was ranked 102nd in the country based on average total risk. To better understand the air toxic problem and promote actions to reduce the risks caused by these pollutants, the Philadelphia Air Toxic Project was initiated by EPA Region III and Air Management Services to develop a more accurate emission inventory, develop modeling systems, identify sources, identify stakeholders and gather background information so a process



can be developed to reduce emissions. Activities associated with the river ports and local airports appear to be significant sources of diesel particulate matter.

AMS has determined that there are health risks associated with the concentrations of air toxics measured at the City's air toxic monitoring sites. Annual averages for each of the compounds at each monitoring site were calculated and used to estimate the risk from inhalation exposure to ambient air for cancer and non-cancer health effects.

The risk calculation is based upon the standard methodology used by EPA. The excess lifetime cancer risk for each of the chemical compounds was calculated using unit risk factors (URFs). The URF is the measure of the probability of developing cancer from exposure over a lifetime to a specified concentration of a given chemical. Air toxics that are measured in Philadelphia that show an excess lifetime cancer risk of 1 or more out of a million are:

1,3-butadiene (Cas RN 106-99-0) - A colorless, non-corrosive gas with a mild aromatic or gasoline-like odor, used primarily as a monomer to manufacture many different types of polymers and copolymers.

acetaldehyde (Cas RN 75-07-0) - A colorless liquid or gas with a fruity odor. It is used to manufacture many other chemicals.

benzene (Cas RN 71-43-2) - A colorless liquid with a pleasant odor. It is used mainly in making other chemicals and plastics, as a solvent, and is found in trace amounts of gasoline.

carbon tetrachloride (Cas RN 56-23-5) - A colorless liquid with an ether-like odor. It is used as a solvent and in making fire extinguishers, refrigerants, and aerosols.

tetrachloroethylene (Cas RN 127-18-4) - A clear liquid with a sweet, chloroform-like odor. It is used in dry cleaning and metal degreasing. Its other common name is perchloroethylene.

The measured ambient concentrations of these toxic pollutants through 2008 are shown by the graphs on the following pages (Figures 12.1 – 16.1). Below each graph is a thematic map developed from the 2002 NATA of the estimated level of risk (Figures 12.2 – 16.2), and a pie chart which shows where emissions come from (Figures 12.3 – 16.3). **Major** sources are large stationary sources such as refineries, power generating stations, and facilities with large boilers. **Area** sources are small stationary sources not included in the major source inventory such as dry cleaners and auto body shops. **On-road** sources are automobiles, trucks, etc. that operate on roads and highways. **Non-road** sources are equipment such as lawn & garden, construction, and airport support. If zero emissions are identified from major, area, on-road, and non-road sources, it is assumed all emissions are from background sources.



Figure 12.1 - 1,3-Butadiene Average Concentration for Five Monitoring Sites




Figure 12.2 - 2002 NATA Total Cancer Risk for 1,3-Butadiene




Figure 12.3 - 1,3-Butadiene Sources of Emissions






Figure 13.1 - Acetaldehyde Average Concentration for Five Monitoring Sites




Figure 13.2 - 2002 NATA Total Cancer Risk for Acetaldehyde




Figure 13.3 - Acetaldehyde Sources of Emissions






Figure 14.1 - Benzene Average Concentration for Five Monitoring Site




Figure 14.2 - 2002 NATA Total Cancer Risk for Benzene




Figure 14.3 - Benzene Sources of Emissions






Figure 15.1 - Carbon Tetrachloride Average Concentration for Five Monitoring Sites




Figure 15.2 - 2002 NATA Total Cancer Risk for Carbon Tetrachloride




Figure 15.3 - Carbon Tetrachloride Sources of Emissions






Figure 16.1 - Tetrachloroethylene Average Concentration for Five Monitoring Sites




Figure 16.2 - 2002 NATA Total Cancer Risk for Tetrachloroethylene




Figure 16.3 - Tetrachloroethylene Sources of Emissions







Figure 17.1 shows the total cancer risk for Philadelphia based on the 2002 NATA data.

Figure 17.1 - 2002 EPA NATA Total Cancer Risk for Philadelphia



The NATA study indicates emissions in Philadelphia for HAPs came mostly from mobile sources (on-road and non-road) as seen in the pie chart in Figure 17.2.

Figure 17.2 - Sum of HAPs Sources of Emissions







Figure 18 shows the cancer risk for the toxics that AMS monitors as seen on the previous pages. It displays the likely rate of risk among a million people over a lifetime per microgram per cubic meter of a toxic.

Figure 18 - Cancer Risks for the Top Five Toxics Compounds





Appendix A: Consequences of Air Pollution



Health Effects

Air pollution contributes to health problems such as asthma, lung disease, and respiratory tract infections. It also can aggravate cardiovascular disease. This concern is greatest in sensitive populations, especially those with lung disorders, young children, and the elderly.

Acid Rain

Acid rain occurs when sulfur dioxide and nitrogen oxides are released into the air, and combine with rain, snow or fog. Acid rain's effects include harming fish, plants, animals, and crops, and eroding building surfaces and national monuments. The effects of acid rain can be offset by reducing the amount of sulfur dioxide released into the air.

Visibility

Haze is caused when particles and gases in the atmosphere scatter or absorb light. The same particles that affect our health also limit our ability to see our surroundings. This affects our quality of living and the beauty of the City of Philadelphia by obscuring many of the national treasures and landmarks that we value.

Climate Change

Global warming refers to an increase in the Earth's temperature, which has the effect of causing climate change. The emission of certain pollutants into the atmosphere which absorb heat energy have sped this process along. Some of the expected long-term changes are a rise in sea level, damage to coastal areas, a variation in precipitation, and other local climate changes. These changes have the potential for altering forests, crop yields, wild life, and water supplies. In 2008, the Mayor's Office of Sustainability, was charged with improving energy efficiency and reducing operating costs in City-owned facilities.



Appendix B: What AMS is Doing to Reduce Air Pollution


- AMS monitors the air for pollutants at many stations throughout the City.
- AMS requires operators of facilities that install or modify sources of air pollution to obtain an air permit from AMS prior to installation. A permit is a list of requirements that includes emission limits and work practice standards as well as testing, monitoring, recordkeeping, and reporting requirements. Sometimes AMS will perform modeling as part of the permit review to assess the impact of new equipment on air quality. A list of recent permit applications received by AMS can be found on our website at www.phila.gov/health/units/ams/index.html. In addition, permits are available to the public. You may contact AMS for information on access to these documents.
- AMS inventories stationary sources of emissions such as factories and other businesses to obtain a current, comprehensive listing of air pollutant emissions for a specific time interval.
- AMS inspects facilities that may cause air pollution or create an air pollution nuisance, meets with and assists facility personnel to achieve compliance, investigates noise and vibration nuisance sources, and responds to citizen complaints and requests for information.
- AMS issues Notices of Violation (NOVs) for sources that are not in compliance, assesses and collects penalties in response to NOVs, initiates orders to abate sources of air pollution, negotiates compliance schedules and agreements to achieve compliance, and refers cases to the Law Department for additional legal remedies including injunctive actions, court orders and consent decrees.
- AMS provides information on economic incentives for cleaning up pollution.
- AMS conducts industry workshops on permitting and emission inventory submittals, provides assistance and training to owners and operators of auto body shops on reducing emissions, and trains asbestos contractors on workplace standards. AMS also provides outreach to educate the public about air quality. Staff attend community fairs, speak at meetings, and visit schools throughout the City.
- AMS is also active in non-regulatory (voluntary) programs. The Philadelphia Diesel Difference is a program that educates owners and operators of diesel engine vehicles about clean diesel technologies such as alternative fuels, anti-idling equipment and diesel engine retrofit devices.



Appendix C: Protecting Yourself and the Environment

If you care about your health, and the health of your neighbors and loved ones, you should also care about the health of the environment. What you do everyday can have a significant impact on the air.

- Avoid overfilling, or “topping off” your car’s gas tank.
- Keep your car's engine tuned up and maintain proper tire pressure.
- When you need to drive, plan ahead so you don't make extra trips.
- Avoid high speeds - fuel efficiency decreases significantly at speeds over 55 mph.
- Drive smoothly to save gas and reduce vehicle emissions.
- Avoid lengthy idling - idling wastes gas.
- Don't rev the engine - this also wastes gas.
- Minimize drag - heavy tow loads, and even driving with the windows open, can create “drag” which reduces fuel efficiency.
- Keep tires properly inflated - under inflated tires decrease gas mileage and shorten tire life. Check the tire pressure in all four tires every two weeks.
- Avoid rough roads and potholes. They are hard on tire and wheel alignment and can also reduce fuel efficiency.



- Use alternative forms of transportation whenever possible, such as carpooling, biking, mass transit, or walking.
- Avoid oil-based paints. Latex paints are much friendlier to the environment, and usually work just as well.

- Conserve energy.
- Don't overheat or overcool your home.
- Turn off lights and appliances when not in use.
- Wash clothes and dishes in full loads.
- Choose Energy Star appliances whenever possible.
- Recycle.
- Report air pollution violations when you become aware of them.
- Take part/respond to hearings/public notices for cleaning up air pollution.
- Learn about local air quality efforts and issues and consider becoming involved in a group that addresses these issues.
- Let your government representatives know that you care about the quality of our City's air.
- Consider switching to wind power or using green building technologies like high-efficiency lighting.





Appendix D: Frequently Asked Questions About Air Quality

What is a criteria pollutant?

A criteria pollutant is one of the six pollutants that are regulated under standards provided by the US Environmental Protection Agency (EPA) to protect the public health and welfare. The criteria pollutants are: ozone, carbon monoxide, nitrogen dioxide, sulfur dioxide, particulate matter, and lead.

Where and how is air quality measured?

The city's air is sampled at stations located throughout Philadelphia. Every minute, monitoring instruments at these stations record the air quality. The data is uploaded to the department's computer system at the AMS Laboratory located at 1501 E. Lycoming Street every 15-minutes. This is called a "real time" system, because the measurements show pollution levels as they are occurring, not after the fact. It allows AMS to evaluate air quality almost continuously. This constant monitoring ensures that no episodes of high pollution are missed and that there are no pollution risks to the public's health.

What does nonattainment mean?

Nonattainment is a designation given to an area that persistently exceeds the ambient air quality standards set for a criteria pollutant. The final designation for nonattainment is given by the EPA after it reviews the recommendations of the State's governor and looks at air quality data for an area. A designation of nonattainment obligates the state or local air agency to identify the causes of pollution, create and implement a strategy that will improve air quality to the point that it meets the standard.

What is an air toxic?

Air toxics or toxic air pollutants, commonly referred to as Hazardous Air Pollutants (HAPs), are air pollutants that cause or may cause cancer or other serious health effects, such as reproductive effects or birth defects, or adverse environmental and ecological effects. The Federal Clean Air Act Amendments of 1990 list 187 pollutants or chemical groups as HAPs.



How do I report an air pollution problem?

For Philadelphia, call 215-685-7580 during AMS office hours from 8 AM to 4:30 PM on weekdays. At all other times, call 311. By calling immediately at the time of the problem, AMS will have a better chance of observing the problem directly and developing an appropriate and timely response.

What information should I report for air pollution problems?

The date, time, location, responsible party (if known) and a description of the problem is requested. A callback name and telephone number is helpful in the event that we are unable to identify the problem. Anonymous calls are also accepted.

What do I do if there's carbon monoxide in my house?

Seek fresh air and dial 9-1-1 for rescue if someone is unconscious or needs help. Otherwise, contact 311. The operator will assist in contacting AMS and other responders as needed.

What should I do if there is unhealthy air quality?

Consider limiting strenuous activity outdoors, especially if you are a member of a sensitive group (the elderly, children, and those with heart or lung problems). Limit the use of your car during daylight hours, and avoid using lawn or garden equipment that requires gasoline.

Why is the ozone layer in the atmosphere considered good, and ozone at ground level considered a health risk?

High in the atmosphere, ozone provides a protective covering for the earth from the sun's ultraviolet (UV) rays, which are harmful. However, ozone low to the ground is formed when certain chemicals and sunlight interact, and is the chief ingredient in smog. It is a strong irritant to the upper respiratory system and eyes, and can cause damage to crops.



Has the air quality in Philadelphia been improving over the years?

Overall Philadelphia's air quality is good and improving. Since 1979, while fluctuating from year to year, the trend of unhealthy and very unhealthy days has steadily declined, especially since 1988. In 1999, the 1997 national standard for 8-hour ozone was incorporated into the AQI, and in 2004, the 1997 national PM_{2.5} standard was added. In 2006, the PM_{2.5} standard became more stringent, and in 2008, the 8-hour ozone standard became more stringent. Because these are more stringent standards, some days that in prior years would have been reported as good fell into the moderate range. This does not reflect a deterioration in air quality.

How was Philadelphia's air quality in 2008?

In 2008, Philadelphia's air quality was rated "good" on 204 days, "moderate" on 144 days, and "unhealthy" on 18 days. These ratings are based on the Air Quality Index (AQI), a system used by cities throughout the country to describe the quality of the air. Air Management Services (AMS) monitors the air quality and enforces the air pollution regulations in the city.

How do I find out more about the air quality in Philadelphia?

The Air Quality Index (AQI) tells you how clean the air is and whether it will affect your health. Philadelphia's Real-Time Air Quality Website is located at www.phila.gov/aqi and provides the most up-to-date information about the air quality in Philadelphia. It lets you know what you should do to protect your health if the air quality is unhealthy. To obtain general air quality information for Philadelphia, arrange for a school or group based presentation or to request a speaker or information booth for a health or environmental fair, contact Air Management Services at 215-685-7586.



Appendix E: Websites

www.aafa.org - Asthma and Allergy Foundation of America

www.airnow.gov/index.cfm?action=airnow.showlocal&CityID=185 - The AQI (Air Quality Index) tells you how clean the air is and whether it will affect your health. Through AIRNow, EPA, NOAA, NPS, state, and local agencies work together to report current and forecast conditions for ozone and particle pollution..

www.airqualitypartnership.org - Ground Level Ozone and Particle Pollution Forecasts

www.atsdr.cdc.gov - Agency for Toxic Substances and Disease Registry: Public health statements on specific toxics and the effects of exposure.

www.cleanair.org - Clean Air Council

www.cleanair.org/dieseldifference - Philadelphia Diesel Difference

www.dep.state.pa.us - Pennsylvania Department of Environmental Protection

www.dieselforum.org - Diesel Technology Forum

www.dvgbc.org - Delaware Valley Green Building Council

www.epa.gov - U.S. Environmental Protection Agency

www.epa.gov/air/data/index.html - AirData presents annual summaries of air pollution data from two EPA databases: the AQS (Air Quality System) database provides air monitoring data - ambient concentrations of criteria air pollutants at monitoring sites, primarily in cities and towns and the NEI (National Emission Inventory) database provides estimates of annual emissions of criteria and hazardous air pollutants from all types of sources.

www.epa.gov/echo - Use ECHO (Enforcement & Compliance History Online) to determine whether compliance inspections have been conducted by EPA or state/local government, violations were detected, enforcement actions were taken, and penalties were assessed in response to environmental law violations.

www.howstuffworks.com - For simple to read answers to a wide variety of science questions, including air pollution, acid rain, and ozone.

www.lungusa.org - American Lung Association website: Information on lung health, air pollution, and related matters.

www.phila.gov/health/units/ams - Philadelphia Department of Public Health, Air Management Services

www.phila.gov/aqi - Philadelphia's Air Quality Website, provides the most up-to-date information about the air quality in Philadelphia and lets you know what you should do to protect your health if the air quality is unhealthy. The tool used to show this is the Air Quality Index (AQI) developed by the Environmental Protection Agency (EPA).

www.scorecard.org - Detailed information on toxics



City of Philadelphia
Department of Public Health
Air Management Services

Appendix F: Contacts

Philadelphia Department of Public Health, Air Management Services	215-685-7580
During non-business hours, operators take calls 8 am to 8 pm on	311
Weekdays, 9 am to 5 pm on Saturdays, and it is closed on Sundays.	
U.S. Environmental Protection Agency, Region III	215-814-5000
Pennsylvania Department of Environmental Protection	484-250-5900
Asthma and Allergy Foundation of America	1-800-476-5536
Clean Air Council	215-567-4004